



Convex bodies with minimal volume product in \mathbb{R}^2 – a new proof

Youjiang Lin*, Gangsong Leng

Department of Mathematics, Shanghai University, Shanghai, 200444, People's Republic of China

ARTICLE INFO

Article history:

Received 6 March 2010

Received in revised form 2 July 2010

Accepted 13 July 2010

Available online 5 August 2010

Keywords:

Convex body

Polar body

Mahler conjecture

Polytope

ABSTRACT

A new proof of the Mahler conjecture in \mathbb{R}^2 is given. In order to prove the result, we introduce a new method – the *vertex removal method*; i.e., for any origin-symmetric polygon P , there exists a linear image ϕP contained in the unit disk B^2 , and there exist three contiguous vertices of ϕP lying on the boundary of B^2 . We can show that the volume-product of P decreases when we remove the middle vertex of the three vertices.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

If $K \subset \mathbb{R}^n$ is an origin-symmetric convex body, let K^* denote its polar body which is defined by

$$K^* = \{x \in \mathbb{R}^n : x \cdot y \leq 1, \forall y \in K\}.$$

Define the volume-product $\mathcal{P}(K)$ of K as

$$\mathcal{P}(K) = \text{Vol}(K)\text{Vol}(K^*).$$

The famous Mahler conjecture [6] is to find a lower bound of $\mathcal{P}(K)$. Is it true that we always have

$$\mathcal{P}(K) \geq \mathcal{P}(B_\infty^n), \quad (1.1)$$

where $B_\infty^n = \{x \in \mathbb{R}^n : |x_i| \leq 1, 1 \leq i \leq n\}$?

For $n = 2$, Mahler [5] proved that the answer is affirmative, and in 1986 Reisner [9] showed that equality holds only for parallelograms. For $n = 2$, a new proof of inequality (1.1) was obtained by Campi and Gronchi [2]. In the n -dimensional case, the conjecture has been verified for some special classes of bodies, namely, for 1-unconditional bodies, [7,10,11], and for zonoids, [3,9].

In 1987, Bourgain and Milman [1] proved that there exists a universal constant $c > 0$ such that $\mathcal{P}(K) \geq c^n \mathcal{P}(B_\infty^n)$, which is now known as the Bourgain–Milman inequality. Very recently, Kuperberg [4] found a beautiful new approach to the Bourgain–Milman inequality. What's especially remarkable about Kuperberg's inequality is that it provides an explicit value for c .

For $n = 2$, Mahler [5] proved that the volume-product of a polygon is concave down as a pair of opposite sides are pivoted and the volume-product can be maximized as all pairs of sides are repeatedly pivoted, at the same time the polygon converges to a regular polygon. In this paper, to prove the Mahler conjecture for $n = 2$, we provide a new method – the *vertex removal method*, which is illustrated as follows: Firstly, we prove that any origin-symmetric polygon P can be transformed

* Corresponding author.

E-mail addresses: linyujiajiang@shu.edu.cn (Y. Lin), gleng@staff.shu.edu.cn (G. Leng).

into a new polygon contained in the unit disk B^2 , in which there are three contiguous vertices lying on the boundary of B^2 (see Theorem 3.1). Next, we prove that the volume-product of the new polygon decreases as the middle vertex of the three contiguous vertices moves toward its adjacent vertices on the boundary of B^2 (see Theorem 3.3). Therefore, we obtain that the volume-product of P decreases when we remove the middle vertex of the three vertices. Compared with Mahler's proof, the *vertex removal method* provides a specific downward path to a square from any origin-symmetric polygon. For $n = 3$, the conjecture could probably be solved following the same idea.

It is worth mentioning that Meyer [8] proved the Mahler conjecture for the general convex bodies in \mathbb{R}^2 . He showed that the volume-product of a convex body is always bigger than that of a triangle and established the case of equality.

2. Definition, notation and preliminaries

As usual, S^{n-1} denotes the unit sphere, B^n the unit ball centered at the origin, o the origin and $\|\cdot\|$ the norm in Euclidean n -space \mathbb{R}^n . If $x, y \in \mathbb{R}^n$, then $x \cdot y$ is the inner product of x and y .

If K is a set, ∂K is its boundary, $\text{int } K$ is its interior, and $\text{conv } K$ denotes its convex hull. Let $\mathbb{R}^n \setminus K$ denote the complement of K ; i.e.,

$$\mathbb{R}^n \setminus K = \{x \in \mathbb{R}^n : x \notin K\}. \quad (2.1)$$

If K is an n -dimensional convex subset of \mathbb{R}^n , then $V(K)$ denotes $\text{Vol}_n(K)$.

Let \mathcal{K}^n denote the set of convex bodies (compact, convex subsets with non-empty interiors) in \mathbb{R}^n . Let \mathcal{K}_o^n denote the subset of \mathcal{K}^n that contains the origin in its interior. Let $h(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$, denote the support function of $K \in \mathcal{K}_o^n$; i.e.,

$$h(K, u) = \max\{u \cdot x \mid x \in K\}, \quad u \in S^{n-1}. \quad (2.2)$$

Let $\rho(K, \cdot) : S^{n-1} \rightarrow \mathbb{R}$, denote the radial function of $K \in \mathcal{K}_o^n$; i.e.,

$$\rho(K, u) = \max\{\lambda \geq 0 \mid \lambda u \in K\}, \quad u \in S^{n-1}. \quad (2.3)$$

A linear transformation (or affine transformation) of \mathbb{R}^n is a map ϕ from \mathbb{R}^n to itself such that $\phi x = Ax$ (or $\phi x = Ax + t$, respectively), where A is an $n \times n$ matrix and $t \in \mathbb{R}^n$. For a convex body P and a linear transformation ϕ , we define ϕP as the linear image of P about ϕ .

Geometrically, an affine transformation in Euclidean space is one that preserves:

- (i) The collinearity relation between points; i.e., three points which lie on a line continue to be collinear after the transformation.
- (ii) Ratios of distances along a line; i.e., for distinct collinear points P_1, P_2, P_3 , the ratio $|P_2 - P_1|/|P_3 - P_2|$ is preserved.

If $K \in \mathcal{K}_o^n$, it is easy to verify that (see p. 44 in [12])

$$h(K^*, u) = \frac{1}{\rho(K, u)} \quad \text{and} \quad \rho(K^*, u) = \frac{1}{h(K, u)}. \quad (2.4)$$

If P is a polygon; i.e., $P = \text{conv}\{A_1, \dots, A_m\}$, where A_i ($i = 1, \dots, m$) are vertices of P . Let a_i denote the vector of A_i . By the definition of the polar body, we have

$$P^* = \bigcap_{i=1}^m \{x \in \mathbb{R}^2 : x \cdot a_i \leq 1\}, \quad (2.5)$$

which implies that P^* is the intersection of m closed half-planes with exterior normal vector a_i . The distance of the straight line $\{x \in \mathbb{R}^2 : x \cdot a_i = 1\}$ from the origin is $1/\|a_i\|$. Thus, if P is an inscribed polygon in a unit circle, then P^* is a polygon circumscribed around the unit circle. In the proof of Theorem 3.3, we shall use these properties.

For $K, L \in \mathcal{K}^n$, the Hausdorff distance is defined by

$$d(K, L) = \min\{\lambda \geq 0 : K \subset L + \lambda B^n, L \subset K + \lambda B^n\}, \quad (2.6)$$

which can be conveniently defined by (see p. 53 in [12])

$$d(K, L) = \max_{u \in S^{n-1}} |h(K, u) - h(L, u)|. \quad (2.7)$$

Therefore, a sequence of convex bodies K_i converges to K if and only if the sequence of support functions $h(K_i, \cdot)$ converges uniformly to $h(K, \cdot)$.

The following lemmas are well-known and important for our proof:

Lemma 2.1. For any origin-symmetric convex body $K \subset \mathbb{R}^n$, $\mathcal{P}(K)$ is linear invariant, that is, for every linear transformation $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$, we have $\mathcal{P}(\phi K) = \mathcal{P}(K)$.

Lemma 2.2. The volume-product $\mathcal{P}(K)$ is continuous under the Hausdorff metric.




Fig. 3.1. Linear transformation ϕ such that $A' = \phi A$.

3. Main result and its proof

Theorem 3.1. In \mathbb{R}^2 , for any origin-symmetric polygon P , there exists a linear image $P' = \phi P$ which satisfies that $P' \subset B^2$ and there exist three contiguous vertices of P' contained in ∂B^2 .

To prove Theorem 3.1, we need the following lemma.

Lemma 3.2. In Fig. 3.1, $A(x_1, y_1)$ and $A'(x_2, y_2)$ are on ∂B^2 . ϕ is a linear transformation from \mathbb{R}^2 to itself such that $A' = \phi A$. If $B(x, y) \in B^2$ and $0 < x_2 < x_1 < x$, then $\phi B \in B^2$.

Proof. Let $B' = \phi B$ and $B' = (x', y')$. Since ϕ is a linear transformation, we get

$$\frac{x}{x_1} = \frac{x'}{x_2} \quad \text{and} \quad \frac{y}{y_1} = \frac{y'}{y_2}.$$

Therefore,

$$x' = \frac{xx_2}{x_1} \quad \text{and} \quad y' = \frac{yy_2}{y_1}.$$

Noting that $x_1^2 + y_1^2 = 1$, $x_2^2 + y_2^2 = 1$ and $x^2 + y^2 \leq 1$, we have

$$\begin{aligned} x'^2 + y'^2 &= \frac{x^2 x_2^2}{x_1^2} + \frac{y^2 y_2^2}{y_1^2} \\ &\leq \frac{x^2 x_2^2}{x_1^2} + \frac{(1 - x^2)(1 - x_2^2)}{1 - x_1^2} \\ &= 1 - \frac{(x_1^2 - x_2^2)(x^2 - x_1^2)}{x_1^2(1 - x_1^2)} \\ &\leq 1. \end{aligned} \tag{3.1}$$

Hence, $\phi B \in B^2$. \square

Now we prove Theorem 3.1.

Proof. Since P is an origin-symmetric polygon, the number of sides of P is even and the opposite sides of P are parallel. Let $A_1, \dots, A_n, B_1, \dots, B_n$ denote all vertices of P , where B_i is the symmetric point of A_i about the origin. Our proof is in three steps.

The first step, transform the parallelogram $A_1 A_2 B_1 B_2$ into the rectangular $A'_1 A'_2 B'_1 B'_2$ inscribed in B^2 . Now P is transformed into P_1 (see (2) or (2)' in Figs. 3.2.1 and 3.2.2).




Fig. 3.2.1. Transforms (1) into (2) or (2)'.




Fig. 3.2.2. Transforms (2) or (2)' into (3).

The second step, transform P_1 into P_2 (see (3) in Fig. 3.2.2). Consider the following two cases for the polygon P_1 ,

- (i) If there exist some vertices of P_1 satisfying

$$\{A'_i : i \in I \subset \{3, \dots, n\}\} \subset \mathbb{R}^2 \setminus B^2,$$

then we transform P_1 into $P_2 \subset B^2$. The transformation shortens the segment $A'_1 A'_2$ into $A''_1 A''_2$ and satisfies that some vertices $\{A'_i : i \in I_1 \subset \{3, \dots, n\}\}$ lie on ∂B^2 .

- (ii) If vertices A'_i and B'_i ($i = 3, \dots, n$) of P_1 satisfy

$$\{A'_3, \dots, A'_n, B'_3, \dots, B'_n\} \subset \text{int } B^2,$$

then we transform P_1 into $P_2 \subset B^2$. The transformation lengthens the segment $A'_1 A'_2$ into $A''_1 A''_2$ and satisfies that some vertices $\{A'_i : i \in I_1 \subset \{3, \dots, n\}\}$ lie on ∂B^2 .

The third step, transform P_2 into P' (see Fig. 3.3). If A''_1, A''_2, A''_i ($i \in I_1$) are three contiguous vertices on ∂B^2 , then this theorem has already been proved; otherwise we rotate P_2 about the origin, we can get a new polygon P'_3 such that $A''_2 A''_i$ parallels the X-axis (see (4) in Fig. 3.3). Then lengthen segments $A''_2 B''_2$ and $A''_i B''_2$ into $A^{(3)}_2 B^{(3)}_2$ and $A^{(3)}_i B^{(3)}_2$, respectively, satisfying that some vertices $\{A^{(3)}_j : j \in I_2 \subset \{3, \dots, i-1\}\}$ lie on ∂B^2 and $\{A^{(3)}_j : j \in \{3, \dots, i-1\} \setminus I_2\} \subset B^2$. By Lemma 3.2, vertices $\{A^{(3)}_{i+1}, \dots, A^{(3)}_n, B^{(3)}_1\}$ are still in the internal of B^2 (see (5) in Fig. 3.3). Let P_3 denote the new polygon. We get $P_3 \subset B^2$.

There are $i-3$ vertices between the vertex $A^{(3)}_2$ and vertex $A^{(3)}_i$. If $i-3 = 1$, then P_3 is the polygon satisfying the theorem. If $i-3 \geq 2$, consider the following two cases:

- (i) If $j-2 \geq i-j$ (where $j \in I_2$), rotate P_3 about the origin such that $A^{(3)}_2 A^{(3)}_j$ parallels the X-axis.
(ii) If $j-2 \leq i-j$ (where $j \in I_2$), rotate P_3 about the origin such that $A^{(3)}_j A^{(3)}_i$ parallels the X-axis.





Fig. 3.3. Transforms (4) into (5).

Fig. 3.4. Polygon P' and its polar body.

We denote the new polygon as P_4 . It is clear that there are less than $i - 3$ vertices between $A_2^{(3)}$ and $A_j^{(3)}$ (or between $A_j^{(3)}$ and $A_i^{(3)}$). By induction, we can get a new polygon $P' \subset B^2$ with three contiguous vertices contained in ∂B^2 after finite transformations. \square

By the above theorem, we consider the volume-product of the polygon with three contiguous vertices contained in ∂B^2 .

Theorem 3.3. Suppose that $P' \subset B^2$ is an origin-symmetric polygon and A, C, B are three contiguous vertices of P' contained in ∂B^2 , then $\mathcal{P}(P'') \leq \mathcal{P}(P')$, where P'' is a new polygon obtained from P' by removing vertices C and C' .

Proof. Suppose that the side AB parallels the X -axis (see Fig. 3.4.) and straight lines l, l_1 and l_2 are three tangent lines to the unit circle B^2 passing through points C, A and B , respectively. Let $A = (-x_0, y_0)$, then $B = (x_0, y_0)$, where $x_0^2 + y_0^2 = 1$. Let $\theta = \angle XOC$. It is clear that $\pi/2 \leq \theta \leq \pi - \arctan(y_0/x_0)$ when the point C is in the second quadrant. We have the following equations of straight lines:

$$l_1 : y - y_0 = \frac{x_0}{y_0}(x + x_0),$$

$$l_2 : y - y_0 = -\frac{x_0}{y_0}(x - x_0),$$

$$l: y - \sin \theta = -\frac{\cos \theta}{\sin \theta}(x - \cos \theta).$$

Let the point N denote the intersection of l and the Y -axis and the point M denote the intersection of l_1 and the Y -axis. We can easily get $N(0, 1/\sin \theta)$ and $M(0, 1/y_0)$. Let H and L denote the intersections of l and l_1, l_2 , respectively. Solve the following systems of equations:

$$\begin{cases} y - \sin \theta = -\frac{\cos \theta}{\sin \theta}(x - \cos \theta) \\ y - y_0 = \frac{x_0}{y_0}(x + x_0) \end{cases} \quad (3.2)$$

and

$$\begin{cases} y - \sin \theta = -\frac{\cos \theta}{\sin \theta}(x - \cos \theta) \\ y - y_0 = -\frac{x_0}{y_0}(x - x_0). \end{cases} \quad (3.3)$$

We get the abscissas of points H and L :

$$x_1 = \frac{y_0 - \sin \theta}{y_0 \cos \theta + x_0 \sin \theta}$$

and

$$x_2 = \frac{y_0 - \sin \theta}{y_0 \cos \theta - x_0 \sin \theta}.$$

Let $S_{\triangle MHL}$ denote the area of $\triangle MHL$, it follows that

$$S_{\triangle MHL} = \frac{x_0}{y_0} \cdot \frac{\sin \theta - y_0}{\sin \theta + y_0}.$$

Let $V = V(P'')$ and $V^0 = V(P''^*)$, where P'' denotes the new polygon obtained from P' by removing vertices C and C' , then $\mathcal{P}(P')$ is a function $f(\theta)$, where

$$f(\theta) = (V + 2x_0(\sin \theta - y_0)) \left(V^0 - \frac{2x_0}{y_0} \cdot \frac{\sin \theta - y_0}{\sin \theta + y_0} \right) \quad (3.4)$$

and

$$\frac{\pi}{2} \leq \theta \leq \pi - \arctan \left(\frac{y_0}{x_0} \right).$$

We have

$$f'(\theta) = 2x_0 \cos \theta \cdot \frac{(V^0 y_0 - 2x_0)(\sin \theta + y_0)^2 + 2y_0(4x_0 y_0 - V)}{y_0(\sin \theta + y_0)^2}. \quad (3.5)$$

In (3.4), since $\cos \theta \leq 0$ and $y_0(\sin \theta + y_0)^2 \geq 0$, to prove $f'(\theta) \leq 0$, let $t = \sin \theta$, we need only to show that $g(t) \geq 0$, where

$$g(t) = (V^0 y_0 - 2x_0)(t + y_0)^2 + 2y_0(4x_0 y_0 - V), \quad t \in [y_0, 1]. \quad (3.6)$$

Let $S_{\square ABA'B'}$ and $S_{\triangle ABM}$ denote the area of a rectangle $ABA'B'$ and triangle ABM , respectively. Since

$$V(P''^*) \geq V(\text{conv}\{A, M, B, A', M', B'\}),$$

we get

$$V^0 \geq S_{\square ABA'B'} + 2S_{\triangle ABM} = 4x_0 y_0 + 2x_0 \left(\frac{1}{y_0} - y_0 \right),$$

therefore,

$$\begin{aligned} V^0 y_0 - 2x_0 &\geq \left(4x_0 y_0 + 2x_0 \left(\frac{1}{y_0} - y_0 \right) \right) y_0 - 2x_0 \\ &= 2x_0 y_0^2 \\ &> 0. \end{aligned} \quad (3.7)$$

Thus, the graph of $g(t)$ is a parabola opening upward. Since the axis of symmetry of the parabola is $t = -y_0$, $g(t)$ is increasing for $t \in [y_0, 1]$. To prove $g(t) \geq 0$, it suffices to prove

$$g(y_0) = 2y_0(2V_0^0 y_0^2 - V) \geq 0. \quad (3.8)$$

Let \mathcal{D} denote the area of a circular segment enclosed by the arc BA' and the chord BA' , then

$$\begin{aligned} V^0 &\geq S_{\square ABA'B'} + 2S_{\triangle ABM} + 2\mathcal{D} \\ &= 4x_0y_0 + 2x_0\left(\frac{1}{y_0} - y_0\right) + 2\mathcal{D} \end{aligned} \quad (3.9)$$

and

$$\begin{aligned} V &\leq S_{\square ABA'B'} + 2\mathcal{D} \\ &= 4x_0y_0 + 2\mathcal{D}. \end{aligned} \quad (3.10)$$

To prove (3.8), it suffices to prove

$$2\left(4x_0y_0 + 2x_0\left(\frac{1}{y_0} - y_0\right) + 2\mathcal{D}\right)y_0^2 \geq 4x_0y_0 + 2\mathcal{D}, \quad (3.11)$$

which is equivalent to

$$2x_0y_0^3 \geq \mathcal{D}(1 - 2y_0^2). \quad (3.12)$$

And since

$$\mathcal{D} \leq (1 - x_0) \cdot 2y_0, \quad (3.13)$$

hence, it suffices to prove

$$x_0y_0^3 \geq y_0(1 - x_0)(1 - 2y_0^2), \quad (3.14)$$

which is equivalent to

$$x_0^3 - 2x_0^2 + 1 \geq 0, \quad (3.15)$$

which is clearly correct when $0 < x_0 < 1$.

We get that $f(\theta)$ is decreasing when $\theta \in [\pi/2, \pi - \arctan(y_0/x_0)]$, hence the function $f(\theta)$ has a minimum value at $\theta = \pi - \arctan(y_0/x_0)$. The point C coincides with the point A when $\theta = \pi - \arctan(y_0/x_0)$. Therefore $\mathcal{P}(P'') \leq \mathcal{P}(P')$. \square

By Theorem 3.3, we can get the following Corollaries 3.4 and 3.5.

Corollary 3.4. If $P \subset \mathbb{R}^2$ is an origin-symmetric polygon, then $\mathcal{P}(P) \geq \mathcal{P}(S)$, where S is a square.

Proof. By Theorems 3.1 and 3.3 and the linear invariance of $\mathcal{P}(P)$, if the number of sides of the polygon P is $2n$, there exists a polygon P_1 with $2(n-1)$ sides satisfying $\mathcal{P}(P_1) \leq \mathcal{P}(P)$. Repeating this process $n-2$ times, we can get a square S satisfying $\mathcal{P}(P) \geq \mathcal{P}(S)$. \square

Corollary 3.5. If $K \subset \mathbb{R}^2$ is an origin-symmetric convex body and $S \subset \mathbb{R}^2$ is an origin-symmetric square, then $\mathcal{P}(K) \geq \mathcal{P}(S)$.

Proof. For any origin-symmetric convex body $K \subset \mathbb{R}^2$, there exists a sequence of origin-symmetric polytopes $\{P_i\}$ converging to K under the Hausdorff metric (see p. 54 in [12]). By Corollary 3.4 and Lemma 2.2., we have

$$\mathcal{P}(K) = \lim_{n \rightarrow \infty} \mathcal{P}(P_i) \geq \mathcal{P}(S). \quad \square \quad (3.16)$$

Acknowledgements

The authors are most grateful to the referee for his many excellent suggestions for improving the original manuscript. The authors would like to acknowledge the support from the National Natural Science Foundation of China (10971128) and Shanghai Leading Academic Discipline Project (S30104).

References

- [1] J. Bourgain, V.D. Milman, New volume ratio properties for convex symmetric bodies in \mathbb{R}^n , *Invent. Math.* 88 (1987) 319–340.
- [2] S. Campi, P. Gronchi, Volume inequalities for L_p -zonotopes, *Mathematika* 53 (2006) 71–80.
- [3] Y. Gordon, M. Meyer, S. Reisner, Zonoids with minimal volume-product – a new proof, *Proc. Amer. Math. Soc.* 104 (1988) 273–276.

- [4] G. Kuperberg, From the Mahler conjecture to Gauss linking integrals, *Geom. Funct. Anal.* 18 (2008) 870–892.
- [5] K. Mahler, Ein minimalproblem für konvexe polygone, *Mathematica (Zutphen)* B 7 (1939) 118–127.
- [6] K. Mahler, Ein Übertragungsprinzip für konvexe Körper, *Časopis Pyest. Mat. Fys.* 68 (1939) 93–102.
- [7] M. Meyer, Une caractérisation volumique de certains espaces normes de dimension finie, *Israel J. Math.* 55 (1986) 317–326.
- [8] M. Meyer, Convex bodies with minimal volume product in \mathbb{R}^2 , *Monatsh. Math.* 112 (1991) 297–301.
- [9] S. Reisner, Zonoids with minimal volume-product, *Math. Z.* 192 (1986) 339–346.
- [10] S. Reisner, Minimal volume product in Banach spaces with a 1-unconditional basis, *J. Lond. Math. Soc.* 36 (1987) 126–136.
- [11] J. Saint Raymond, Sur le volume des corps convexes symétriques, *Seminaire d'initiation al Analyse*, 1980–1981, *Publ. Math. Univ. Pierre et Marie Curie*, Paris, 1981, pp. 1–25.
- [12] R. Schneider, *Convex Bodies: The Brunn–Minkowski Theory*, in: *Encyclopedia Math. Appl.*, vol. 44, Cambridge University Press, Cambridge, 1993.